

Prediction of Dragon's Blood Tree (*Dracaena cinnabari* Balf.) stand sample density on Soqotra Island (key study)

Irena Hubálková

Mendel University in Brno, Faculty of Forestry and Wood Technology, Department of Forest Botany, Dendrology and Geobiocoenology, Zemědělská 3, 613 00 Brno, Czech Republic
irena.hubalkova@mendelu.cz

Abstract

Dracaena cinnabari Balf. On The Soqotra Island is a spectacular relict of the Tethys tropical forest. This unique endemic plant producing medicinally valuable sap covered larger area in the past. There is no natural regeneration of this species except inaccessible localities with steep slopes. All seedlings are threatened by goats grazing. Age structure of *Dracaena* populations indicates maturity and overmaturity depending on browsing.

The objective of this work is to predict growing dynamic of Dragon's Blood Trees in permanent sample plot at Firmihin, where is the largest closed stand of *Dracaena* species. The prediction and visualization of variation in the number of trees during 100 years is based on realized direct field measurements supported by mathematical calculations. The study presents options in forest regeneration together with caution about the threats in the way of its implementation.

Key words - *Dracaena cinnabari*, Soqotra, stand visualization, trees density prediction

INTRODUCTION

The Soqotra Archipelago (see Fig. 1) is located in the northwestern part of the Indian Ocean. It consists from two main islands Soqotra (also written Sokotra, Socotra, Suqutra) and Abdalkuri (Abd El Kuri), and the smaller ones, Samhah and Darsa, called The Brothers, and of the rocks of Cal Farun and Hertha (Mies and Beyhl, 1996). Politically, this archipelago belongs to the Republic of Yemen. With an area of 3625 km² is Soqotra the largest island in the Arab world (Elie, 2002), lies about 235 km east of the Horn of Africa (12°18'-12°42' N latitude and 53°18'-54°32' E longitude) and the highest elevations is reached in Haghier Mountains on Soqotra (over 1500 m a. s. l.).




Fig. 1 – Geographical location of Soqatra Island.

Many authors have provided information about climate on Soqatra, but this information seems to be based on individual experience from short-term stays on island or considered from wider climatic characteristics from Indian Ocean or nearby mainlands (Habrová, 2007). According to Mies and Beyhl (1996), the islands are situated in the arid tropical zone where evapotranspiration generally surpasses precipitation by far. The climate of the ecoregion is influenced by the southwest (summer) and northeast (winter) monsoons. The south-west monsoon (from May to September) brings only humidity, the north-east monsoon (from November to March) is milder but brings expected winter rain (Fleitmann et al., 2007) and the rest of the year can be expected only dry weather conditions.

Tab. I - Annual statistics of 'daily mean values' of selected climatic variables calculated from completed values in 'ideal year' (2000-2004) at Firmihin locality (Král, 2005).

Variable	Min	Max	Mean
Air temperature [°C]	19.3	28.9	23.7
Air humidity [%]	36.6	99.5	72.6
Wind speed [m/s]	0.3	8.2	2.6
Precipitation [mm/day]	0.0	207.5	1.1

The Soqatra Island is undoubtedly a most precious natural asset, not only for the Republic of Yemen, but for many people from around the world (Yucer, 1988). In 2003, the archipelago was declared the first UNESCO Man and Biosphere Reserve in Yemen, because of Soqatra Island was recognised as one of the best preserved island ecosystems on Earth (Miller et al., 2006). Habrová et al. (2009) mentioned island ranking into the World Heritage Site in 2008. Separated from continent during the Tertiary period the island hosts no indigenous mammals whatsoever, but its floral endemism rate making

it one of the most biodiverse island in the world (Grant, 2005). 825 species of flowering plants and ferns were recorded on the archipelago, from which 12 genera and 307 species (37%) are thought to be endemic (Miller et al., 2006). According to Buček et al. (2004), also the majority of woody species forming natural forest, woodland and/or shrub communities ranks among endemic species. Important of them are arborescent frankincense trees (*Boswellia* sp.), myrrh trees (*Commiphora* sp.), arborescent spurges (*Euphorbia* sp.), a rare endemic shrub *Dirachma socotrana* and the only known wild pome granate species (*Punica protopunica*). Typical for Soqotra are remarkable succulent woody species particularly the endemic cucumber tree (*Dendrosycios socotrana*), desert rose (*Adenium obesum*, ssp. *socotranum*) and endemic *Dorstenia gigas*. The unique vegetation formation is the evergreen woodland dominated by the famous Dragon's Blood Tree (*Dracaena cinnabari*), Soqotra's most iconic plant (Miller et al., 2006). Soqotra Island is rightly compared with Mauritius, the Galapagos or the Canary Islands. It is given by floristic richness and level of endemism. Also fauna of Soqotra is rich in species and not fully investigated yet.

For thousands of years, the people of Soqotra have lived with and used the biological resources in a sustainable manner (Mies and Beyhl, 1996) harvesting only what they needed and practising rotational grazing (Miller et al., 2006). However, the Soqotra Archipelago with an estimated human population ranging from 40 to 80000 people is described as one of the poorest and most disadvantaged group of islands anywhere in the world (Elie, 2002). Most of people are concentrated in the capital Hadiboh and in its surroundings. They speak Soqotri and Arabic. The people have survived through fishing, ranging of livestock (mostly goats) and home gardening, particularly date palms (Ceccolini, 2000). Since antiquity, the island was famous for its Dragon's Blood, obtained from the sap of *Dracaena cinnabari*, (Mies and Beyhl, 1996). Nowadays, the island is losing its isolation rapidly. This is due to industrial and tourism development.

The genus *Dracaena* comprises between 60 and 100 species and recent taxonomic ambiguity has caused its classification within three families i.e. *Agavaceae*, *Liliaceae* and *Dracaenaceae*, the latter as a family arching over the former confusion (Adolt and Pavliš, 2004). Representatives of the *Dracaena* genus have survived in woodlands on dry margins of the Tethys tropical forest since the Tertiary Period. Most of the *Dracaena* species grow as shrubs or geophytes often having ornamental potential (Adolt and Pavliš, 2004). There are some species having the growth habit of a tree: *Dracaena cinnabari* Balf. on Soqotra, *Dracaena serrulata* Baker in Southwestern Arabia, *Dracaena ombet* Kotschy & Peyr in Eastern Africa, *Dracaena schizantha* Baker on Macaronesian Islands, *Dracaena draco* L.

in Moricci, *Dracaena tamaranae* A. Marrero, R.S. Almeida and M. González-Martín in Canary Islands and *Dracaena steudneri* Engl. in Ethiopia and Eastern Africa (Bekele, 2007).

Dracaena cinnabari is an evergreen tree with a typical umbrella-shaped crown due to a "dracoid" ramification of branches (Adolt and Pavliš, 2004). *Dracaena* species are exceptional among monocotyledonous plants because of their capacity for secondary thickening of stems and roots (Habrová et al., 2009). Its area of distribution ranges from an altitude of 150 m to 1600 m above sea level, it dominates above 600 m a. s. l. (Petroncini, 2001). Dragon's Blood Tree is not widely spread over the Soqotra Island. It is fixed to the area of the large central plateau of Diksam, the central granite massive of Haggeher and the eastern area of Hamadero, Sirahon and Kilisan (Petroncini, 2001). Plant density is not homogenous. Analyses show that the area of *Dracaena* woodland land-cover class on Soqotra reaches 3658 ha, i. e. 1.1 % of the total island area (Král and Pavliš, 2006). There is no natural seeding except of inaccessible places with steep slopes.

Multiyear activities of a Mendel University team from Brno has brought important data concerning particularly ecophysiology, phenology, morphology, growth characteristics, estimation of age, population dynamics of *Dracaena cinnabari*, site conditions or species composition. Precise geobiocoenological differentiation was created for recognizing the detailed state of natural conditions of the island (Buček, 2003). Mendel University team also installed a weather station including an automatic data logger at Firmihin in November 2000, at an approximate altitude of 440 m above sea level. According to Habrová (2007), it is clear, that only a long-term, continuous period of measurement at different locations on the island can lead to an understanding of its present climate. That is why other 5 mini-stations measuring only air temperature and air humidity were placed at different localities in 2004 (one of these 5 mini-stations has been stolen on Skant just after one month).

According to mentioned researches, the population of *Dracaena cinnabari* is getting old. Stands density decreases. There is absence of natural regeneration at most sites of natural range. The species are threatened mainly by omnipresent goat grazing, extraction of blood-red sap and fuelwood production.

MATERIAL AND METHODS

Data collection and field measurement

Data about Soqotra Archipelago and *Dracaena cinnabari* were collected. It was focused on general information about Soqotra Island (localization, climatic conditions, natural environment and nature conservation, local people and their natural resources utilization). Other part of introduction dealt with *Dracaena* genus and Soqotra's most iconic plant *Dracaena cinnabari* (habit, distribution, utilization and its importance). Also the connections with causes of threats to this species were mentioned.

Field measurements were realized at Firmihin, where the largest closed stand of Dragon's Blood Trees was identified. One square sample plot with side length 100 m in locality of the youngest *Dracaena* trees was chosen. The inventory of each tree in sample plot was carried out. Measuring-tape for girth measuring in 1.3 m and crown diameter, hypsometer Silva, camera and field notebook were used. Usage of laser rangefinder was expected, but the instrument was confiscated by soldiers at the airport in capital of Yemen. All trees were photographed. The central point of sample plot was located by GPS Trimble-Juno SB and stabilized by metal spile.

Data processing

Collected data were compiled in Microsoft Office Program Excel, copied to electronic scratch pad and applicated to Stand Visualization System (SVS). The scratch pad files were saved with the ending tbl (xls ending was useless). The data applicable to SVS consist of a simple stand table containing species, diameter at breast height (DBH) in cm, height in m, ratio (= crown height/tree height in m), crown radius in m, X and Y coordinates. Except these parameters 'sausage-shape' sections of branches (see Fig. 3) and circumferential inflorescence were counted during fieldwork. Also columns Tree class and Crown class were added in Microsoft Office Program Excel, which are necessary to processing in SVS.

In general, 100 tables for 100 year prediction of this sample plot were generated. Prediction of trees sample density was developed on basis of previous long-term measurement and statistical evaluation made by members of Mendel University team (Adolt and Pavliš, 2004) at Firmihin.

Next step consisted in assignment of all trees into 16 age categories by Habrová (2005) (Tab. III). The categories were called from DR16 (seed) to DR1 (the oldest trees). The number of trees in particular category provided the vector of a population (Habrová et al., 2009). The age of one branch section reaches about 19 years (Adolt and Pavliš, 2004). It was necessary to predefine tree habits with different species codes in Tree Designer of SVS (see Fig. 2). The auxiliary view in SVS Tree Designer was used to display trees as they appear in an SVS image. Every created tree form was marked with assigned species code, which allows user to make changes to the form definition parameters. The tree class list shows all the tree class values currently represented for the current species. The crown class list shows all crown class values currently represented for the current species and current tree class. It was necessary to create a new plant form for *Dracaena* sp. based on an existing form. The form usable as the model was selected and then the species, tree class and crown class for the new plant form were modified.

According to SVS Manual created by McGaughey (2002), SVS generates images depicting stand conditions represented by a list of individual stand components (e. g. trees, shrubs) using detailed geometric models. The images produced by SVS provide a readily understood representation of stand conditions.




Fig. 2 – Modeling of tree age classes in SVS Tree Designer.

In order to construct a model of population development in the following years, it was necessary to compile a generally applicable matrix created by Habrová et al. (2009). The matrix (Tab. II) is constructed of values for the probability of dying out within a given age category and the value of the average age of one section of a branch. The values along a diagonal give the probability that a plant will persist in the same category during one year. The matrix was multiplied by a particular vector in a numerical computing environment MATLAB (i.e., by the actual number of trees in particular categories). Graph expressing the decrease in the number of trees during 100 years was created in MATLAB by mentioned multiplying the matrix and vectors.

Tab. II - Matrix and the vector (= real number of trees fallen into each category for a given "number of narrowed branch sections").

	1st flower	1 - 2	3 - 4	5 - 6	7 - 8	9 - 10	11 - 12	13 - 14	15 - 16	17 - 18	19 - 20	21 - 22	23 - 24	25 <	Vector
1st flower	0,98824	0	0	0	0	0	0	0	0	0	0	0	0	0	4
1 - 2	0,01176	0,97368	0	0	0	0	0	0	0	0	0	0	0	0	2
3 - 4	0	0,02632	0,97178	0	0	0	0	0	0	0	0	0	0	0	8
5 - 6	0	0	0,02626	0,97206	0	0	0	0	0	0	0	0	0	0	16
7 - 8	0	0	0	0,02627	0,97125	0	0	0	0	0	0	0	0	0	22
9 - 10	0	0	0	0	0,02625	0,97251	0	0	0	0	0	0	0	0	17
11 - 12	0	0	0	0	0	0,02628	0,96975	0	0	0	0	0	0	0	19
13 - 14	0	0	0	0	0	0	0,02621	0,96822	0	0	0	0	0	0	16
15 - 16	0	0	0	0	0	0	0	0,02617	0,96801	0	0	0	0	0	4
17 - 18	0	0	0	0	0	0	0	0	0,02616	0,96619	0	0	0	0	4
19 - 20	0	0	0	0	0	0	0	0	0	0,02611	0,96354	0	0	0	0
21 - 22	0	0	0	0	0	0	0	0	0	0	0,02604	0,94713	0	0	0
23 - 24	0	0	0	0	0	0	0	0	0	0	0	0,02560	0,94270	0	1
25 <	0	0	0	0	0	0	0	0	0	0	0	0	0,02548	0,95833	1

RESULTS

The research was realized on Firmihin, located in the central part of Soqotra Island. The dense *Dracaena* woodland found at Firmihin is unique in the world. Exact location of the stabilized central point of the sample plot is X 175964,9968856; Y 1381889,1487503; Z 544,8 (WGS-84 reference system).

114 *Dracaena* trees were identified on selected sample plot. In average trees have had from 9 to 10 branch sections (see Fig. 3).

Tab. III - Number of trees belonging to each age category.

Marking of age category	Number of branch sections	Number of trees
DR16	seed	0
DR15	seedling	0
DR14	the first flowering plant	4
DR13	1 and 2	2
DR12	3 and 4	8
DR11	5 and 6	16
DR10	7 and 8	22
DR9	9 and 10	17
DR8	11 and 12	19
DR7	13 and 14	16
DR6	15 and 16	4
DR5	17 and 18	4
DR4	19 and 20	0
DR3	21 and 22	0
DR2	23 and 24	1
DR1	25 <	1



Fig. 3 – Branches segregated by narrowed 'sausage-shape' sections (photo by authoress).

As mentioned above, 100 tables for 100 year prediction of this sample plot were generated in Microsoft Office Program Excel. The data in first table (age class, DBH, height, ratio, crown radius, X a Y coordinates) showed present condition of the trees (in the year 2010). The data in hundredth table showed the same parameters 100 years later (in the year 2110). The diameter at breast height (DBH) has been averagely increasing 0.053 cm per year and crown radius 0.0089 m per year (Adolt and Pavliš, 2004).

First and last table were copied from Microsoft Office Program Excel to electronic scratch pad (Tab. IV), which is form applicable in Stand Visualization System (SVS).

Tab. IV Parameters of *Dracaena* trees in 2010 displayed in electronic scratch pad.

	Soubor	Úpravy	Formát	Zobrazení	Nápověda										
DR8	38.22	5.1	0.24	1.32	1	98	99	1	5	5	0	0	0	0	0
DR7	41.4	5.4	0.37	1.84	1	98	99	1	13	8	0	0	0	0	0
DR8	31.21	5.71	0.32	1.36	1	98	99	1	20	2	0	0	0	0	0
DR8	32.17	4.38	0.4	1.53	1	98	99	1	26	6	0	0	0	0	0
DR11	27.39	5.31	0.18	1.01	1	98	99	1	37	10	0	0	0	0	0
DR10	19.15	5.33	0.26	1.62	1	98	99	1	2	14	0	0	0	0	0
DR5	52.23	4.81	0.49	2.31	1	98	99	1	10	18	0	0	0	0	0
DR12	24.52	4.88	0.18	0.89	1	98	99	1	4	12	0	0	0	0	0
DR10	34.08	5.97	0.44	1.27	1	98	99	1	17	11	0	0	0	0	0
DR12	33.44	6.91	0.77	1.53	1	98	99	1	20	15	0	0	0	0	0
DR7	47.77	5.13	0.37	2.06	1	98	99	1	28	13	0	0	0	0	0
DR10	38.85	5.46	0.4	1.65	1	98	99	1	44	9	0	0	0	0	0
DR5	48.41	5.56	0.4	2.17	1	98	99	1	48	5	0	0	0	0	0
DR6	46.82	5.12	0.39	2.36	1	98	99	1	46	17	0	0	0	0	0
DR10	35.03	6.38	0.25	1.43	1	98	99	1	49	19	0	0	0	0	0
DR8	22.61	4.5	0.2	0.68	1	98	99	1	50	50	0	0	0	0	0
DR14	35.03	5.22	0.41	1.86	1	98	99	1	40	23	0	0	0	0	0
DR5	36.62	5.96	0.42	1.84	1	98	99	1	39	26	0	0	0	0	0
DR7	28.66	6.37	0.38	1.19	1	98	99	1	30	20	0	0	0	0	0
DR9	28.82	5.94	0.3	1.26	1	98	99	1	24	25	0	0	0	0	0
DR10	40.45	5.14	0.38	1.86	1	98	99	1	15	27	0	0	0	0	0
DR10	29.3	5.15	0.29	1.33	1	98	99	1	11	21	0	0	0	0	0
DR10	28.03	4.91	0.62	0.9	1	98	99	1	3	29	0	0	0	0	0
DR14	45.22	5.31	0.35	1.88	1	98	99	1	6	35	0	0	0	0	0
DR7	38.22	5.09	0.35	1.29	1	98	99	1	17	40	0	0	0	0	0
DR11	28.03	5.02	0.37	1.11	1	98	99	1	22	32	0	0	0	0	0

Graph (see Fig. 4) expresses prediction of tree density between 2010 and 2110. The graphical representation shows, that from 114 trees, approximately 73 trees will remain. Thus the number of trees will decrease by 36 %.




Fig. 4 – Graph expressing the decrease in the number of *Dracaena* trees during 100 years; ten years interval between 0 and 100 is represented.

Two models were created in SVS on the basis of proven results. The first model shows present condition of the trees (see Fig 5). The second model expresses the decrease in the number of trees during 100 years (see Fig. 6). The disappearance of the trees was chosen in consideration of age, vitality and degree of damage. The most frequent type of damage was stem scar caused by extraction of blood-red sap. An intensive pasture of goats is expected. That is the reason why no natural regeneration and/or artificial plantation are included.




Fig. 5 – Model in SVS showing *Dracaena* trees sample density in 2010.




Fig. 6 – Model in SVS showing *Dracaena* trees sample density in 2110.

It is necessary to mention, that this prediction is very optimistic, because currently (in 2010) there are the youngest *Dracaena* trees in Firmihin on chosen sample plot. Natural disasters or other unexpected events are not assumed.

DISCUSSION

Age determination of monocotyledons in general, is very difficult task (Habrová et al., 2009). One study (Adolt and Pavliš, 2004) on the estimation of the age of *Dracaena* tree species has been published. According to mentioned study, background is an indirect method reflecting relationship between the number of flowering periods and the actual age of a specimen. The relation can be well defined through specific branching of arborescent *Dracaena* species characterized by markedly swollen branches segregated individually by narrowed sections or 'sausage-shaped' sections. Based on this method, one branch section reaches about 19 years and *Dracaena* achieves an age up to 650 years. But even so it is hard to determine the age of stem before crown branching, because there are no growth rings in cross section of the stem. Habrová (2005) mentioned that there are three main growth

stages in *Dracaena*'s ontogeny. The seedling creates no trunk some time after germination, just a single rosette is growing and the number of leaves is increasing. After some time the trunk is creating, but there is still only one single rosette on the top (Attorre et al., 2007). The branching and crown forming is related to the maturing stage (certain stem height) and ability to flower. Height growth cessation happens with ability to flower and moves between 40 and 1200 cm (Habrová, 2005).

According to the method developed by Adolt and Pavliš (2004), the crown of the oldest tree on sample plot was 532 years old (28 branch sections). There were no seedlings without stem and only three young trees creating stem with no branching.

Based on statistical analyses as well as on direct field observations, *Dracaena* populations on Soqotra do not regenerate to a great extent and their age structure generally indicates overmaturity (Král and Pavliš, 2006). This research results confirm the claim by Adolt and Pavliš (2004), that with 95 % probability, it is possible to expect that larger part of the world's most extensive *Dracaena* woodland complex at Firmihin will be in the stage of intensive disintegration within 30 to 77 years. Graphical representation as well as two models created by SVS show absence of natural regeneration. There are mostly middle aged trees in a sample plot, despite the fact that it is locality with one of the youngest trees at Firmihin. The results demonstrate gradual degradation and disappearance of trees.

Some scientists (Attorre et al., 2007) attribute the Dragon's Blood Tree reduction to climatic changes. According to Attorre et al. (2007), the predicted climate change may lead to a 45 % loss of *Dracaena* potential distribution area by 2080. The study published by Attorre et al. (2007) suggests that the original distribution of *Dracaena cinnabari* has been significantly reduced in the past. They hypothesise that a combination of factors may have contributed to its reduction (human activities, soil erosion, increased aridity and biotic interactions). They also suppose that current pattern of distribution, though fragmented may be primarily explained in terms of response to climatic constraints. Attorre et al. (2007) opine that Dragon's Blood Tree is able to colonise suitable areas if the present climatic conditions will remain stable and if a reduction of grazing and human pressure will occur.

Mendel University team does not consider climate change as a significant factor in the process of *Dracaena* tree reduction. We are tending to opinion that grazing by livestock, predominantly by goats, is the reason for its decline. Cattle grazing and excessive consumption of fuelwood and building timber most markedly represent

an increasing anthropogenic pressure on natural resources of Soqotra (Buček, 2003). Seed production capability of Dragon's Blood Trees is optimal. However, thanks to intensive browsing, there is the absence of natural regeneration except inaccessible steep slopes. The only way to prevent the decrease in stand density is a strict reserve and protection from grazing at Firmihin. Mentioned solution appears unrealistic in conditions of Soqotra Island, where most of people support livestock grazing. Even so, the only way to prevent the degradation and disappearance of trees is to prevent grazing damage.

The *Dracaena* woodland on Soqotra Island is rightly considered as one of the oldest forest community on Earth. It is a unique phytocoenose, which covered larger area in the past. Mendel University team activities lead towards promotion of *Dracaena cinnabari* regeneration, which count tree nursery establishment, endorsement and plants protection. The team is attempting to plant cultivated plants on the original localities of the species. Obstacles to successful behaviour consist partially in distrust local people to foreigners, habitual way of traditional grazing management, long periods of dry weather conditions etc. It is necessary to support optimizing of Dragon's Blood Tree age structure.

SUMMARY

The aim of this study was to predict and visualize variation in the number of Dragon's Blood Trees during 100 years in permanent sample plot at Firmihin on the Island of Soqotra. Field measurements were realized in Firmihin, where the largest closed stand of *Dracaena cinnabari* was identified. One square sample plot of side length 100 m in locality of the youngest Dragon's Blood Trees was chosen. The inventory of each tree in sample plot was carried out. Collected data were compiled in Microsoft Office Program Excel, copied to electronic scratch pad and applicated to Stand Visualization System (SVS). The prediction of growing dynamic was developed on basis of previous long-term measurement and statistical evaluation made by Mendel University team. The mortality of trees during next 100 years was evaluated, graphically figured out and the models of stand development were created on the basis of matrix indicating the probability of tree within branch section category persistence (Habrová et al., 2009).

114 *Dracaena* trees on selected sample plot were identified. According to the method based on architectural age and statistical analyses published by Adolt and Pavliš (2004), the crown of the oldest tree was 532 years old (28 branch sections). The average trees have about 9 sections. There were found no seedlings without stem and only three young trees

with created stem without branching. The graphical representation created in MATLAB shows, that from 114 trees, approximately 73 trees will remain 100 years later. Thus the number of trees will decrease by 36 %.

Two models were created in SVS on the basis of proven results. The first model shows trees sample density in 2010. The second model expresses the decrease in the number of 41 trees during 100 years. The curve (see Fig. 4) as well as two models created by SVS show absence of natural regeneration. Most likely this is due to livestock grazing, primarily by goats. The only way to prevent the degradation and disappearance of trees is to prevent grazing damage.

ACKNOWLEDGEMENT

This study was supported by IGA 12/2010, Faculty of Forestry and Wood Technology, Mendel University in Brno, Czech Republic.

REFERENCES

ADOLT R., PAVLIŠ, J., 2004: Age structure and growth of *Dracaena cinnabari* populations on Socotra. In: Trees Structure and Function. Vol. 18, N. 1: 43–53 p.

ATTORE, F., FRANCESCONI, F., TALEB, N., SCHOLTE, P., SAED, A., ALFO, M., BRUNO, F., 2007: Will dragonblood survive the next period of climate change? Current and future potential distribution of *Dracaena cinnabari* (Socotra, Yemen). In: Biological Conversation. Vol. 138, N. 3–4: 430–439 p.

BEKELE, T. A., 2007: Useful Trees of Ethiopia: Identification, propagation and management in 17 Agroecological Zones. In: RELMA in ICRAF Project, Nairobi: 552 p.

BUČEK, A., 2003: Geobiocoenological Typology and Agroforestry as a Tool for Sustainable Development of Soqotra Island. Aden University Printing & Publishing house, Aden: 97–108 p.

BUČEK, A., HABROVÁ, H., KRÁL, K. 2004: Geobiocenologická typologie ostrova Sokotry a její aplikace při tvorbě ekologické sítě. In: Fyzickogeografický sborník 2 - Kulturní krajina. Masarykova univerzita, Brno: 26–42 p. ISBN 80-210-3597-8.

CECCOLINI, L., 2000: The home gardens of Soqotra. Florence: Department of Plant Biology-Laboratory, University of Florence, [cit. 2010-06-05]. Available on WWW: < http://www.etfrn.org/ETFRN/newsletter/news36/nl36_oip7.html >.

ELIE, S. D., 2002: A historical genealogy of Socotra as an object of mythical speculation, scientific research and development experiment. Brighton: The Centre for Culture, Development and Environment, University of Sussex, [cit. 2010-02-25]. Available on WWW: <<http://www.aiys.org/webdate/socot.html>>.

FLEITMANN, D., BURNS, S. J., MANGINI, A., MUDELSEE, M., KRAMERS, J., VILLA, I., NEFF, U., AL-SUBBARY, A. A., BUETTNER, A., HIPPLER, D., MATTER, A., 2007: Holocene ITCZ and Indian monsoon dynamics recorded in stalagmites from Oman and Yemen (Socotra). In: Quaternary Science Reviews. Vol. 26, N. 1–2: 170–188 p.

GRANT, G., 2005: Socotra: Hub of the Frankincense Trade. In: Explorations: An Undergraduate Research Journal: 119–138, [cit. 2010-07-19]. Available on WWW: <<http://undergraduatestudies.ucdavis.edu/explorations/2005/grant.pdf>>.

GUPTA, D., BLEAKLEY, B., GUPTA, R. K., 2008: Dragon's blood: Botany, chemistry and therapeutic uses. In: Journal of Ethnopharmacology. Vol. 115, N. 3: 361–380 p.

HABROVÁ, H., 2005: Geobiocenologická typologie a agrolesnictví jako nástroj pro tvorbu ekologické sítě ostrova Sokotra (Jemenská republika). Brno: Department of Forest Botany, Dendrology and Geobiocoenology, Mendel University in Brno, Czech Republic. Dissertation.

HABROVÁ, H., 2007: Climate of Soqotra with a View to Rais. In: Tayf – The Soqotra Newsletter. Vol. 4. 15–16 p.

HABROVÁ, H., ČERMÁK, Z., PAVLIŠ, J., 2009: Dragon's blood tree – Threatened by overmaturity, not by extinction: Dynamics of a *Dracaena cinnabari* woodland in the mountains of Soqotra. In: Biological Conservation. Vol. 142, N. 4: 772–778 p.

KRÁL, K., 2005: Assessment and mapping of forest and shrub geobiocoenoses by geoinformation methods. Brno: Department of Forest Botany, Dendrology and Geobiocoenology, Mendel University in Brno, Czech Republic. Dissertation.

KRÁL, K., PAVLIŠ, J., 2006: The first detailed land cover map of Socotra Island by Landsat /ETM+ data. In: International Journal of Remote Sensing. Vol. 27, N. 15: 3239–3250. ISSN 0143-1161.

MCGAUGHEY, R. J., 2002: The Stand Visualization System Manual. USDA Forest Service, Pacific Northwest Research Station, [cit. 2010-05-18]. Available on WWW: <<http://forsys.cfr.washington.edu/winsvs/manual.pdf>>.

MIES, B. A., BEYHL, F. E., 1996: The vegetation ecology of Soqotra. Soqotra – Proceedings of the First International Symposium on Soqotra Island: Present & Future. In: United Nations Publications, New York: 35–82 p.

MILLER, A., MORRIS, M., WRANIK, W., KNEES, S., 2006: Soqotra – Land of the Dragon's Blood Tree. Royal Botanic Garden Edinburgh, Edinburgh: 64 p. ISBN 1-872291-40-6.

PETRONCINI, S., 2001: Survey and monitoring of *Dracaena cinnabari* Balf. Fil. in Soqotra Island (Yemen). Florence: Department of Plant Biology-Laboratory, University of Florence, [cit. 2010-06-08]. Available on WWW:

<http://www.etfrn.org/ETFRN/newsletter/news36/nl36_oip7.html >.

YUCER, O., 1988: Preface. In: Soqotra – Proceedings of the First International Symposium on Soqotra Island: Present & Future. United Nations Publications, New York: 9–10 p.